PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. Keep people and pets out of treated areas until sprays have dried.

FIRST AID:

If In Eves: Hold eve open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eve. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment

For Emergency Medical Information, call the National Pesticide Information Center at 1-800-858-7378

ENVIRONMENTAL HAZARDS: Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

PRODUCT INFORMATION

Treats up to 3.000 Square Feet (Quart Treats up to 6,000 Square Feet [1/2 Gallon] Treats up to 12.000 Square Feet [1 Gallon]

WHERE TO USE:

- Around patios, walkways, driveways and fences.
- In vegetable or flower gardens and around trees and shrubs and other landscaped areas.
- In large areas:
- For lawn renovation or replacement
- To prepare garden plots for planting

AMOUNT TO USE-

For Best Results: 6 fl.oz. (12 Tbs) per gallon of water.

For areas with small or seedling weeds: 3 fl. oz. (6 Tbs) per gallon of water per 300 square feet.

NOTICE: Read the entire label. Use only according to label directions. Before using this product, read "Warranty Disclaimer", "Inherent Risks of Use", and "Limitations of Remedies". If terms are unacceptable, return at once unopened.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

This is an end-use product. The manufacturer does not intend and has not registered it for reformulation.

General Information

This product kills most broadleaf and grass weeds. This product may be used for non-selective weed control around patios, walkways, driveways and fences, in vegetable or flower gardens and around trees and shrubs and other landscaped areas, in large areas for lawn removal prior to replacing turfgrasses, and for weed control in garden plots prior to planting. Applications to plant foliage are rainfast in approximately two hours and afterward will not be affected by rainfall or irrigation

How This Product Works

The active ingredient in this product is active against a substance found only in plants. It enters the plant through foliage and is translocated throughout the plant killing both roots and top growth. This product is not active in soil and c nnot move through the soil to damage desirable plants. Sprays not the role by clants are broken down into natural materials.

What To Expect

Weeds may begin to wilt within a few hours, but complete vill may require 1 to 2 weeks. Small, rapidly ving we as mer be used with a single application while older, mature weer's may equin re-applicatio. If re-application is necessary, wait until new growth appears.

Important Use Precautions

Do not spray during windy conditions or allow the spray to contact desirable plants such as flowers, ground evers or lan areas, as they will be severely damaged or killed. In situations where weeds and desirable plants are in close proximity, use a piece of cardboard or plastic to shield desirable plants from the spray. If a desirable plant is spraved by mistake, immediately rinse the plant with water. Before sprays have dried, avoid inadvertent transfer of the sprav solution from treated plants to desirable plants or tracking spray from treated areas onto lawns or other desirable vegetation.

Do not apply through any type of irrigation system.

Keep people and pets out of treated areas until sprays have dried.

SPRAYER CLEANUP

Rinse spraver and all spraver parts three times with water. Apply rinse waters to bare soil or graveled area away from desirable vegetation. After careful cleaning, spraver may be used for other products.



MIXING INSTRUCTIONS

(Plastic, aluminum or stainless steel):

• Use 6 fl. oz. (12 Tbs) per gallon c way Spot treat or splay evenly over 300 m.

Ortho* Dial in Spray

(Do not ad , water

pr. v ev ally over measured area. ablespoon(Tbs) = 3 teaspoons (tsp) 1 fl. oz. = 2 Tbs = 6 tsp

⁺ This product can react with certain meta surfaces and should not be applied with galvanized or unlined steel sprayers.

APPLICATION

For Best Results

- Apply when weather is warm (above 60°F) and sunny to stimulate systemic movement from foliage to roots.
- To avoid spray drift to desirable vegetation adjust sprav nozzle to produce large droplets and spray when wind is calm.
- Spray foliage until thoroughly wet, but not past the point of runoff.
- Apply when weeds are small and actively arowing, before seeds develop,

Treatment of Vines

To control vines growing on fences, mature trees, or poles, cut the vines back to a height of 3 to 4 feet and spray thoroughly. Avoid spray contact with desirable plants or lawn.

For control of vines climbing on trees or shrubs with green bark, remove vines by cutting off at the base of the plant and spray after the vines have begun to regrow. Shield trees or shrubs with a sheet of cardboard or plastic to prevent contact with spray.

potato, pumpkin, radish, rhubarb, rutabaga, spinach squash, sweet potato, tomatillo, tomato(from seed) turnip, watercress and vams. Lawn grass and the following caneberries may be planted 7 days after application: Blackberry blueberry, boysenberry, currant, dewberry elderberry, gooseberry, huckleberry, loganberry, olallieberry, and raspberry (black and red).

Tomatoes from transplant and unlisted fruits vegetables and herbs may be planted 30 days after application.

Turf Replacement

beco po dry.

Harvesting

harvesting.

Replanting

after application.

Terms and Conditions of Use If terms of the following Warranty Disclaimer Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other use

 Treats up to 12.000 Square Feet [1 Gallon] alvahosatot: N-(ahosahonomothyl) alveino isopropylamino salt 19.0%

Kills Weeds, Roots and All

Rainproof in 2 Hours

giyphosate. N-(phosphononethy) giycine, is	opropyidi i ille sait18.0%
IERT INGREDIENTS:	
DTAL INGREDIENTS	100.00%

Treats up to 3.000 Square Feet [Quart]

Treats up to 6.000 Square Feet [1/2 Gallon]

Contains 1.6 pounds per gallon glyphosate, isopropylamine salt (1.2 pounds per gallon glyphosate acid).

ACTIVE INGREDIENTS:



EPA REG. NO. 10088-107-68562 EPA EST, 10088-WI-1 NET CONTENTS: 1 1 5 Gallon(s)

Distributed By: Total Solutions[™] PO Box 240014 • Milwaukee WI 53224

kill an existing lawn prior to replacement skip one m wing before spraving and apply at the higher rate to fl. oz. (12 Tbs.) per gallon of water. Control wil be letter if the grass is actively growing, so water 2 to 3 days before and after application if the soil

Wait an additional 7 days after the last application before raking, tillage prior to reseeding or sodding.

If used to control weeds around fruit or nut trees caneberries, or grapevines, allow 21 days before

Flowers, shrubs, and trees may be planted 1 day

These fruits, vegetables, and herbs may be planted 3 days after application: Beans, beet greens, broccoli, brussels sprouts, cabbage, carrot cauliflower, celery, chard(Swiss), chicory, collards corn, cucumber, egaplant, endive, garlic, gourds horseradish. Jerusalem artichoke, kale, kohlrabi leek, lentils, lettuce, melons, mustard greens, okra onion, rape greens, parsley, parsnips, peas, pepper

constitutes acceptance of the terms under Warranty Disclaimer. Inherent Risks of Use and Limitations of Remedies

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description of the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER Reapply to any green patches that remain after 7 MAKES NO EXPRESS NOR IMPLIED WARRANTY OF MER-CHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Lack of performance, or other unintended consequences may occur due to such factors as use of the product contrary to label instructions (including adverse conditions noted on the label, such as unfavorable temperature. soil conditions, etc.), abnormal conditions (such as extreme weather conditions), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

STORAGE AND DISPOSAL

Do Not Contaminate Water, Food, or Feed By Storage or Disposal

Pesticide Storage: Store in a dry place no lower in temperature than 50°F or higher than 120°F.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions. contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Nonrefillable container. Do not reuse or refill container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat procedure two more times.