

to contact treated surface for a minimum of 3 minutes prior to wiping. Reapplication is necessary if surface is dried before contact time is achieved. When used on toys or food contact surfaces such as countertops, tables and appliances, thoroughly rinse surfaces with potable water before use. Do not use on utensils, dishes or glasses.

To Sanitize Non-Food Contact Surfaces: After hard, non-porous surface is cleaned, hold can upright 6" to 8" from area to be treated. Spray until surface is thoroughly wet. Allow to dry without wiping or allow spray to contact treated surface for a minimum of 3 minutes prior to wiping. Reapplication is necessary if surface is dried before contact time is achieved.

To Kill Poliovirus Type 1: After surface is cleaned, hold can upright 6" to 8" from area to be treated. Spray until surface is thoroughly wet. Allow to dry without wiping or allow spray to contact treated surface for a minimum of 5 minutes prior to wiping. Reapplication is necessary if surface is dried before contact time is achieved. When used on toys or food contact surfaces such as countertops, tables and appliances, thoroughly rinse surfaces with potable water before use. Do not use on utensils, dishes or glasses.

To Kill (TB) Mycobacterium bovis: Pre-clean surfaces, hold can upright 6" to 8" from area to be treated. Spray until surface is thoroughly wet. Allow to dry without wiping or allow spray to contact treated surface for a minimum of 10 minutes prior to wiping. Reapplication is necessary if surface is dried before contact time is achieved. When used on toys or food contact surfaces such as countertops, tables and appliances, thoroughly rinse surfaces with potable water before use. Do not use on utensils, dishes or glasses.

For use as a fungicide against Trichophyton interdigitale (Athlete's Foot): Apply to clean surface holding can upright 6" to 8" from area to be treated. Spray until surface is thoroughly wet. Allow to dry without wiping or allow spray to contact treated surface for a minimum of ten minutes prior to wiping. Reapplication is necessary if surface is dried before contact time is achieved.

For Spot Sanitization of Soft Non-food Contact Surfaces: After surface is cleaned, hold can upright 6" to 8" from area to be treated. Spray until surface is thoroughly wet. Allow to dry without wiping or allow spray to contact treated surface for a minimum of three minutes prior to wiping. Reapplication is necessary if surface is dried before contact time is achieved.

For Control of Mold and Mildew: Apply by holding can upright 6" to 8" from surface to be treated. Spray until surface is completely wet. Let air dry. Repeat application at weekly intervals or when mildew growth reappears.

***Kills HIV on Pre-Cleaned Environmental Surfaces Previously Soiled with Blood/Body Fluids** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission

of human immunodeficiency virus* Type 1 (HIV) (associated with AIDS): **Special Instructions for Cleaning and Decontamination Against HIV on Surfaces/Objects Soiled with Blood/Body Fluids: Personal Protection:** When handling items soiled with blood or body fluids, use barrier protection: disposable latex gloves, gowns, masks and eye coverings. **Cleaning Procedure:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this disinfectant. **Contact Time:** Spray surface to be treated 3-4 seconds, from a distance of 6-8 inches, until surface is wet. Allow spray to dry without wiping or allow spray to contact treated surface for a minimum of three minutes except for Poliovirus Type 1 which requires five minutes, prior to wiping. **Disposal of Infectious Materials:** Autoclave blood and other body fluids and disposed of according to federal, state and local regulations for infectious waste disposal.

This product is not for use on medical devices or medical equipment surfaces.

Storage and Disposal

Pesticide Storage: Store in a cool, dry (preferably locked) place inaccessible to children and pets and away from direct sunlight, windows, flames and other sources of heat or ignition. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed on site or at an approved waste disposal facility. **Container Disposal:** Non-refillable container. Do not refill or reuse this container. Do not puncture or incinerate! If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions.

PRECAUTIONARY STATEMENTS
Hazards to Humans & Domestic Animals
CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

First Aid
If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes.

If on Skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-866-836-8855 for emergency medical treatment information.

Physical or Chemical Hazards
 Flammable. Contents under pressure. Do not use or store near fire, sparks or heated surfaces. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Incineration of container may cause explosion.

Manufactured by



CLAIRE MANUFACTURING COMPANY
 1000 Integram Drive | Pacific, MO 63064
 800.252.4731
 www.clairemfg.com



MADE IN USA
 with domestic and foreign components.



DISINFECTANT SPRAY Q

Country Fresh Scent

For Use on
**Hard, Non-porous Environmental Surfaces
 and Soft Spot Surface Sanitizer applications.**

Use in:
 Healthcare, Schools, Hotels, Restaurants,
 Commercial Establishments, Homes

**MULTIPURPOSE DISINFECTANT
 KILLS 99.9% GERMS
 CONTROLS MOLD AND MILDEW
 ELIMINATES ODORS**

Bactericidal - Antibacterial - Fungicidal
 *Virucidal - Tuberculocidal



Nonfood Compounds
 Program Listed C1
 Reg #158919

**Keep Out of Reach
 of Children**
CAUTION
 See back panel for First Aid and
 additional Precautionary
 Statements.

Active Ingredients:
 Alkyl (60% C14, 30% C16, 5% C18, 5% C12)
 dimethyl benzyl ammonium chloride..... 0.16%
 Alkyl (68% C12, 32% C14)
 dimethyl ethylbenzyl ammonium chloride ... 0.16%
Other Ingredients 99.68%
 Total 100.00%
 EPA Reg. No. 706-111
 EPA Est. 33595-MO-4
 Refer to bottom of container for production facility

Net Wt. 17 oz. (1 lb. 1 oz.) 482 grams

Disinfects on hard, non-porous surfaces and soft spot surface sanitizer applications. For non-food application use on: High touch surfaces, door knobs, light switches, toilets, sinks, bathtubs, showers, kitchen appliance surfaces, garbage cans. Disinfects food contact surfaces such as tables, counter tops, appliances, and toys in Kitchens, Restaurants, Commercial Establishments and Homes.
 Sanitizes soft non-food contact surfaces as a spot treatment.

Claire Disinfectant Spray Q sanitizes and kills 99.9% of *Staphylococcus aureus* (staph) and *Enterobacter aerogenes* in 3 minutes on pre-cleaned, hard, non-porous surfaces.

Claire Disinfectant Spray Q kills the following bacteria, fungi, and *viruses in 3 minutes on pre-cleaned, hard, non-porous surfaces.

Bacteria: *Pseudomonas aeruginosa* (ATCC 15442), *Salmonella enterica* (ATCC 10708), *Staphylococcus aureus* (ATCC 6538), *Acinetobacter baumannii* Multi Drug Resistant (ATCC 19606), Carbapenem Resistant *Escherichia coli* (CDC 81371), Carbapenem Resistant *Klebsiella pneumoniae* (ATCC BAA-1705), *Enterobacter aerogenes* (ATCC 13048), *Enterococcus faecium* (ATCC 51559), *Escherichia coli* (ATCC 11229), *Klebsiella pneumoniae* (ATCC 4352), *Listeria monocytogenes* (ATCC 19117), *Staphylococcus aureus- MRSA* (ATCC 33592), *Staphylococcus aureus- CA-MRSA Genotype USA 300* (NARSA NRS 384), *Staphylococcus aureus- CA-MRSA Genotype USA 400* (NARSA NRS 123), *Shigella dysenteriae* (ATCC 11835), *Streptococcus pyogenes* (ATCC 19615), *Staphylococcus aureus -VISA* (CDC HIP 5836), *Enterococcus faecalis -VRE* (ATCC 51575), *Campylobacter jejuni* (ATCC 29428), *Candida albicans* (ATCC 10231).

***Viruses:** Adenovirus Type 2*, 2009-H1N1 Influenza A virus*, Influenza B virus*, Human Immunodeficiency virus type 1 (HIV-1)*, Herpes simplex virus type 1*, Herpes simplex virus type 2*, Rotavirus*, Vaccinia virus*, Rhinovirus type 37*.

Kills Poliovirus type 1 in 5 minutes on pre-cleaned hard, non-porous surfaces.

Fungi: *Trichophyton interdigitale* (Athlete's Foot Fungus) (ATCC 9533) (10 minute contact time)

Kills *Mycobacterium bovis* (TB) in 10 minutes at 22.5°C on pre-cleaned hard, non-porous surfaces.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Shake Well Before Each Use. Point valve opening toward area to be cleaned, press down firmly and spray evenly.

To Disinfect: After hard, non-porous surface is cleaned, hold can upright 6" to 8" from area to be treated. Spray until surface is thoroughly wet. Allow to dry without wiping or allow spray



PRODUCT # C-11001